

2004
Update



China

EPIDEMIOLOGICAL FACT SHEETS
ON HIV/AIDS AND SEXUALLY TRANSMITTED INFECTIONS



Joint United Nations Programme on HIV/AIDS

UNAIDS

UNHCR • UNICEF • WFP • UNDP • UNFPA
UNODC • ILO • UNESCO • WHO • WORLD BANK



World Health
Organization

HIV/AIDS estimates

In 2003 and during the first quarter of 2004, UNAIDS and WHO worked closely with national governments and research institutions to recalculate current estimates on people living with HIV/AIDS. These calculations are based on the previously published estimates for 1999 and 2001 and recent trends in HIV/AIDS surveillance in various populations. A methodology developed in collaboration with an international group of experts was used to calculate the new estimates on prevalence and incidence of HIV and AIDS deaths, as well as the number of children infected through mother-to-child transmission of HIV. Different approaches were used to estimate HIV prevalence in countries with low-level, concentrated or generalised epidemics. The current estimates do not claim to be an exact count of infections. Rather, they use a methodology that has thus far proved accurate in producing estimates that give a good indication of the magnitude of the epidemic in individual countries. However, these estimates are constantly being revised as countries improve their surveillance systems and collect more information.

Adults in this report are defined as women and men aged 15 to 49. This age range covers people in their most sexually active years. While the risk of HIV infection obviously continues beyond the age of 50, the vast majority of those who engage in substantial risk behaviours are likely to be infected by this age. The 15 to 49 range was used as the denominator in calculating adult HIV prevalence.

Estimated number of adults and children living with HIV/AIDS, end of 2003

These estimates include all people with HIV infection, whether or not they have developed symptoms of AIDS, alive at the end of 2003:

Adults and children	840,000		
Low estimate	430,000		
High estimate	1,500,000		
Adults (15-49)	830,000	Adult rate (%)	0.1
Low estimate	430,000	Low estimate	0.1
High estimate	1,400,000	High estimate	0.2
Children (0-15)			
Low estimate			
High estimate			
Women (15-49)	190,000		
Low estimate	95,000		
High estimate	320,000		

Estimated number of deaths due to AIDS

Estimated number of adults and children who died of AIDS during 2003:

Deaths in 2003	44,000
Low estimate	21,000
High estimate	75,000

Estimated number of orphans

Estimated number of children who have lost their mother or father or both parents to AIDS and who were alive and under age 17 at the end of 2003:

Current living orphans	
Low estimate	
High estimate	

UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance

Global Surveillance of HIV/AIDS and sexually transmitted infections (STIs) is a joint effort of WHO and UNAIDS. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, initiated in November 1996, guides respective activities. The primary objective of the Working Group is to strengthen national, regional and global structures and networks for improved monitoring and surveillance of HIV/AIDS and STIs. For this purpose, the Working Group collaborates closely with national AIDS programmes and a number of national and international experts and institutions. The goal of this collaboration is to compile the best information available and to improve the quality of data needed for informed decision-making and planning at national, regional, and global levels. The Epidemiological Fact Sheets are one of the products of this close and fruitful collaboration across the globe.

Within this framework, the Fact Sheets collate the most recent country-specific data on HIV/AIDS prevalence and incidence, together with information on behaviours (e.g. casual sex and condom use) which can spur or stem the transmission of HIV.

Not unexpectedly, information on all of the agreed upon indicators was not available for many countries in 2003. However, these updated Fact Sheets do contain a wealth of information which allows identification of strengths in currently existing programmes and comparisons between countries and regions. The Fact Sheets may also be instrumental in identifying potential partners when planning and implementing improved surveillance systems.

The fact sheets can be only as good as information made available to the UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance. Therefore, the Working Group would like to encourage all programme managers as well as national and international experts to communicate additional information to them whenever such information becomes available. The Working Group also welcomes any suggestions for additional indicators or information proven to be useful in national or international decision-making and planning.

Assessment of the epidemiological situation 2004

China is experiencing one of the most rapidly expanding HIV epidemics in the world. According to the national surveillance system, it is estimated that in 2003, there were 840,000 people infected with HIV (range 430,000 to 1.02 million). The first major outbreak started in Yunnan province in 1989 among injecting drug users (IDU). Since 1999, there has been a 30% annual rate of increase of reported HIV infections. Based on this calculation, it is projected that without concerted prevention and treatment efforts, the number people living with HIV/AIDS (PHA) in China will exceed 10 million by 2010, if there is no enough effort to HIV control. To some degree, HIV/AIDS affects all parts of the country, however the bulk of the impact can be found in about 200 of China's 2,800 counties. Most of these heavily affected counties are rural and poor.

Injecting drug use is becoming more common with many IDUs sharing needles; consequently, the introduction of HIV in these groups has resulted in a rapid spread of HIV. Based on sentinel surveillance conducted in 2003, HIV prevalence rate among IDUs was found to range from 44% to 85% in selected communities of drug users in Yunnan and Xinjiang provinces are HIV-infected. The percentage of IDUs who report sharing of equipment increased from 31.7% in 1999, 33.7% in 2000 to 45% in 2001. A second epidemic of HIV was found among fomal plasma donor (FPDs) in provinces in middle of China. According to epidemiological survey, HIV prevalence rate among FPDs was found to range from 5% to 60% in selected communities of FPDs in Henan provinces. Limited studies among the FPDs have revealed HIV prevalence rates of 1-39% in selected communities of Hebei, Hubei, Hunan, Anhui, Shanxi, Shaanxi, and other provinces with less prevalence rates such as Jilin (county and township), Shandong and Sichuan. Anhui, Hubei and Henan are the 3 of most affected provinces with the most people on ARV treatment.

Heterosexually transmitted HIV is occurring primarily in the eastern provinces of China. Heterosexual transmission is fueled by an increasing commercial sex trade and by the large number of migrants moving to these provinces in search of labor. The percentage of female prostitutes who do not use condoms decreased from 66.7% in 1999, 49.1% in 2000 to 37.4% in 2001 (median).

A related problem is a dramatic increase in the prevalence of STDs, particularly ulcerative STDs that are known to facilitate the transmission of HIV. The number of reported STDs has increased more than 100-fold between 1985 and 2001. Another worrisome factor is that many sex workers are also drug users, thereby providing a bridge between the high risk IDU group and the general population. Finally, HIV has been identified in some urban areas among men who have sex with men but this population is stigmatized and is difficult to reach in terms of conducting serologic surveys or prevention outreach.

Basic indicators

For consistency reasons the data used in the table below are taken from official UN publications.

DEMOGRAPHIC DATA	YEAR	ESTIMATE	SOURCE
Total population (thousands)	2004	1,313,309	UN population division database
Female population aged 15-24 (thousands)	2004	102,186	UN population division database
Population aged 15-49 (thousands)	2004	741,994	UN population division database
Annual population growth rate (%)	1992-2002	0.9	UN population division database
% of population in urban areas	2003	38.2	UN population division database
Average annual growth rate of urban population	2000-2005	3.2	UN population division database
Crude birth rate (births per 1,000 pop.)	2004	14.2	UN population division database
Crude death rate (deaths per 1,000 pop.)	2004	7.1	UN population division database
Maternal mortality rate (per 100,000 live births)	2000	56	WHO (WHR2004)/UNICEF
Life expectancy at birth (years)	2002	71.1	World Health Report 2004, WHO
Total fertility rate	2002	1.8	World Health Report 2004, WHO
Infant mortality rate (per 1,000 live births)	2000	31	World Health Report 2004, WHO
Under 5 mortality rate (per 1,000 live births)	2000	37	World Health Report 2004, WHO

SOCIO-ECONOMIC DATA	YEAR	ESTIMATE	SOURCE
Gross national income, ppp, per capita (Int.\$)	2002	4,390	World Bank
Gross domestic product, per capita % growth	2001-2002	7.2	World Bank
Per capita total expenditure on health (Int.\$)	2001	224	World Health Report 2004, WHO
General government expenditure on health as % of total expenditure on health	2001	37.2	World Health Report 2004, WHO
Total adult illiteracy rate	2000	14.8	UNESCO
Adult male illiteracy rate	2000	7.9	UNESCO
Adult female illiteracy rate	2000	22.1	UNESCO
Gross primary school enrolment ratio, male	2000/2001	114	UNESCO
Gross primary school enrolment ratio, female	2000/2001	114	UNESCO
Gross secondary school enrolment ratio, male	2000/2001	77	UNESCO
Gross secondary school enrolment ratio, female	2000/2001	58	UNESCO

Contact address

UNAIDS/WHO Working Group on Global
HIV/AIDS and STI Surveillance
20, Avenue Appia
CH - 1211 Geneva 27
Switzerland
Fax: +41-22-791-4834

email: hivstrategicinfo@who.int or
estimates@unaids.org

website: <http://www.who.int/hiv>
<http://www.unaids.org>

Extracts of the information contained in these fact sheets may be reviewed, reproduced or translated for research or private study but not for sale or for use in conjunction with commercial purposes. Any use of information in these fact sheets should be accompanied by the following acknowledgment "UNAIDS/WHO epidemiological fact sheets on HIV/AIDS and Sexually Transmitted Infections, 2004 Update".

HIV prevalence in different populations

This section contains information about HIV prevalence in different populations. The data reported in the tables below are mainly based on the HIV database maintained by the United States Bureau of the Census where data from different sources, including national reports, scientific publications and international conferences are compiled. To provide a simple overview of the current situation and trends over time, summary data are given by population group, geographical area (Major Urban Areas versus Outside Major Urban Areas), and year of survey. Studies conducted in the same year are aggregated and the median prevalence rates (in percentages) are given for each of the categories. The maximum and minimum prevalence rates observed, as well as the total number of surveys/sentinel sites, are provided with the median, to give an overview of the diversity of HIV-prevalence results in a given population within the country. Data by sentinel site or specific study from which the medians were calculated are printed at the end of this fact sheet.

The differentiation between the two geographical areas Major Urban Areas and Outside Major Urban Areas is not based on strict criteria, such as the number of inhabitants. For most countries, Major Urban Areas were considered to be the capital city and - where applicable - other metropolitan areas with similar socio-economic patterns. The term Outside Major Urban Areas considers that most sentinel sites are not located in strictly rural areas, even if they are located in somewhat rural districts.

HIV sentinel surveillance*


Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
Pregnant women	Major urban areas	N-Sites										1.00	1.00	2.00	2.00	3.00				
		Minimum										0	0	0	0	0				
		Median										0	0	0	0	0				
		Maximum										0	0	0	0	0				
	Outside major urban areas	N-Sites					1.00	1.00	1.00				1.00	9.00	11.00	2.00				
		Minimum					0.20	0	0.04				0	0	0	0				
		Median					0.20	0	0.04				0	0.12	0.22	0.25				
		Maximum					0.20	0	0.04				0	0.47	1.63	0.50				
Sex workers	Major urban areas	N-Sites									15.00	15.00	17.00	19.00	19.00	18.00	1.00			
		Minimum									0	0	0	0	0	0	10.30			
		Median									0	0	0	0	0	0	10.30			
		Maximum									0	0.71	0.70	2.55	6.19	10.29	10.30			
	Outside major urban areas	N-Sites			1.00	1.00	1.00	1.00	1.00	1.00	4.00	4.00	5.00	5.00	5.00	4.00	1.00			
		Minimum			0	0	0.38	0	0	0	0	0	0	0	0	0	0	0.44		
		Median			0	0	0.38	0	0	0	0.18	0	0	0.20	0.40	0.12	0.44			
		Maximum			0	0	0.38	0	0	0	0.50	1.50	2.67	23.08	2.70	2.90	0.44			
Injecting drug users	Major urban areas	N-Sites					2.00	2.00	1.00	1.00	8.00	10.00	10.00	13.00	11.00	16.00				
		Minimum					0	0	0	0	0	0	0	0	0	0				
		Median					0	0	0	0	0	0	0.61	3.85	1.11	0.43				
		Maximum					0	0	0	0	0.68	9.29	27.52	40.45	24.40	20.46				
	Outside major urban areas	N-Sites			2.00	2.00	1.00	9.00	9.00	9.00	11.00	19.00	13.00	20.00	21.00	10.00				
		Minimum			38.81	26.64	3.73	0	0	0	0.15	0	0	0.55	0	0				
		Median			41.98	47.31	3.73	3.73	2.67	1.54	8.27	16.67	22.00	16.40	17.00	1.95				
		Maximum			45.14	67.98	3.73	81.82	56.25	47.13	73.21	76.79	70.27	85.12	81.77	82.49				
STI patients	Major urban areas	N-Sites						1.00	1.00		24.00	22.00	26.00	28.00	70.00	73.00				
		Minimum						0	0		0	0	0	0	0	0				
		Median						0	0		0	0	0	0	0	0				
		Maximum						0	0		0.30	0.45	0.43	0.40	1.54	2.06				
	Outside major urban areas	N-Sites				1.00	1.00	1.00		1.00	5.00	8.00	4.00	16.00	22.00	9.00				
		Minimum				0	0	0		1.40	0	0	0	0	0	0				
		Median				0	0	0		1.40	0.03	0.93	0	0.44	0.44	0				
		Maximum				0	0	0		1.40	6.70	15.26	0	11.27	11.32	0.61				
Men having sex with men																				
Tuberculosis patients																				

*Detailed data by site can be found in the Annex.


Maps & charts

Mapping the geographical distribution of HIV prevalence among different population groups may assist in interpreting both the national coverage of the HIV surveillance system as well in explaining differences in levels of prevalence. The UNAIDS/WHO Working Group on Global HIV/AIDS and STI Surveillance, in collaboration with the WHO Public Health Mapping Team, Communicable Diseases, is producing maps showing the location and HIV prevalence in relation to population density, major urban areas and communication routes. For generalized epidemics, these maps show the location of prevalence of antenatal surveillance sites.

Trends in antenatal sentinel surveillance for higher prevalence countries, or in prevalence among selected populations for countries with concentrated epidemics, are a new addition. These are presented for those countries where sufficient data exist.



Trends in HIV prevalence in high risk groups



Median prevalence and ranges are shown in areas with more than one sentinel site.

The boundaries and names shown and the designations used on the map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement. WHO 2004, all rights reserved.

Reported AIDS cases

Following WHO and UNAIDS recommendations, AIDS case reporting is carried out in most countries. Data from individual AIDS cases are aggregated at the national level and sent to WHO. However, case reports come from surveillance systems of varying quality. Reporting rates vary substantially from country to country and low reporting rates are common in developing countries due to weaknesses in the health care and epidemiological systems. In addition, countries use different AIDS case definitions. A main disadvantage of AIDS case reporting is that it only provides information on transmission patterns and levels of infection approximately 5-10 years in the past, limiting its usefulness for monitoring recent HIV infections.

Despite these caveats, AIDS case reporting remains an important advocacy tool and is useful in estimating the burden of HIV-related morbidity as well as for short-term planning of health care services. AIDS case reports also provide information on the demographic and geographic characteristics of the affected population and on the relative importance of the various exposure risks. In some situations, AIDS reports can be used to estimate earlier HIV infection patterns using back-calculation. AIDS case reports and AIDS deaths have been dramatically reduced in industrialized countries with the introduction of Anti-Retroviral Therapy (ART).

1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
						1	0	2	0	0	2	3	5	23	29	52	38	126	136
1999	2000	2001	2002	2003	Total		UNK		Date of last report										
230	233	231			1111		10/26/2001												

Curable sexually transmitted infections (STIs)

The predominant mode of transmission of both HIV and other STIs is sexual intercourse. Measures for preventing sexual transmission of HIV and STIs are the same, as are the target audiences for interventions. In addition, strong evidence supports several biological mechanisms through which STIs facilitate HIV transmission by increasing both HIV infectiousness and HIV susceptibility. Thus, detection and treatment of individuals with STIs is an important part of an HIV control strategy. In summary, if the incidence/prevalence of STIs is high in a country, then there is the possibility of high rates of sexual transmission of HIV. Monitoring trends in STIs provides valuable insight into the likelihood of the importance of sexual transmission of HIV within a country, and is part of second generation surveillance. These trends also assist in assessing the impact of behavioural interventions, such as delaying sexual debut, reducing the number of sex partners and promoting condom use.

Clinical services offering STI care are an important access point for people at high risk for both STIs and HIV. Identifying people with STIs allows for not only the benefit of treating the STI, but for prevention education, HIV testing, identifying HIV-infected persons in need of care, and partner notification for STIs or HIV infection. Consequently, monitoring different components of STI prevention and control can also provide information on HIV prevention and control activities within a country.

STI syndromes

Reported cases	1996	1997	1998	1999	2000	2001	2002	2003	Incidence 2003
Comments:									
Source:									

Syphilis prevalence, women

Percent of blood samples taken from pregnat women aged 15-49 that test positive for syphilis - positive reaginic and treponemal test-during routine screening at selected antenatal clinics.

Year	Area	Rate	Range
Comments:			
Source:			

Estimated prevalence of curable STIs among female sex workers

- Chlamydia

Year	Area	Rate	Range
2000-2001	Urban	56.6 (rate)	
1997-1999	Urban	32.2 (rate)	

Comments:

Source:

Ministry of Health of the People’s Republic of China, WHO Regional Office for the Western Pacific. Prevalence Survey of Sexually Transmitted Infections among Sex Workers and Truck Drivers in China 1999-2000.

- Gonorrhoea

Year	Area	Rate	Range
2000-2001	Urban	37.8	
1997-1999	Urban	8.8	

Comments:

Source:

Ministry of Health of the People’s Republic of China, WHO Regional Office for the Western Pacific. Prevalence Survey of Sexually Transmitted Infections among Sex Workers and Truck Drivers in China 1999-2000.

Estimated prevalence of curable STIs among female sex workers (continued)

- Syphilis

Year	Area	Rate	Range
1997-1999	Urban	5.4	
2000-2001	Urban	10	

Comments:

Source: van den HA. High prevalence of syphilis and other sexually transmitted diseases among sex workers in China: potential for fast spread of HIV. AIDS 2001, 15:753-759.

- Trichomoniasis

Year	Area	Rate	Range
2000-2001	Urban	43.2	
1997-1999	Urban	12.5	

Comments:

Source: Ministry of Health of the People's Republic of China, WHO Regional Office for the Western Pacific. Prevalence Survey of Sexually Transmitted Infections among Sex Workers and Truck Drivers in China 1999-2000.

Health service and care indicators

HIV prevention strategies depend on the twin efforts of care and support for those living with HIV or AIDS, and targeted prevention for all people at risk or vulnerable to the infection. It is difficult to capture such a large range of activities with one or just a few indicators. However, a set of well-established health care indicators may help to identify general strengths and weaknesses of health systems. Specific indicators, such as access to testing and blood screening for HIV, help to measure the capacity of health services to respond to HIV/AIDS - related issues.

Access to health care

Indicators	Year	Estimate	Source
% of population with access to health services - total			
% of population with access to health services - urban			
% of population with access to health services - rural			
Contraceptive prevalence rate (%)	1998	90.7	UNICEF/UNPOP
Percentage of contraceptive users using condoms			
% of births attended by skilled health personnel	2000	89.3	WHO
% of 1-yr-old children fully immunized - DPT	2002	79	WHO/UNICEF
% of 1-yr-old children fully immunized - Measles	2001	79	WHO/UNICEF
% of ANC clinics where HIV testing is available			

Number of adults (15-49) with advanced HIV infection receiving ARV therapy as of June 2004

Adults on treatment

Number:	7,400
Source:	WHO

Estimated number of adults (15-49) in need of treatment in 2003

Adults needing treatment

Number:	100,000
Source:	WHO/UNAIDS

Coverage of HIV testing and counselling

Number of public and NGO services providing testing and counselling services.

Year	Area	N=
------	------	----

Comments:
Source:

Knowledge and behaviour

In most countries the HIV epidemic is driven by behaviours (e.g.: multiple sexual partners, injecting drug use) that expose individuals to the risk of infection. Information on knowledge and on the level and intensity of risk behaviour related to HIV/AIDS is essential in identifying populations most at risk for HIV infection and in better understanding the dynamics of the epidemic. It is also critical information in assessing changes over time as a result of prevention efforts. One of the main goals of the 2nd generation HIV surveillance systems is the promotion of a standard set of indicators defined in the National Guide (Source: National AIDS Programmes, A Guide to Monitoring and Evaluation, UNAIDS/00.17) and regular behavioural surveys in order to monitor trends in behaviours and to target interventions.

The indicators on knowledge and misconceptions are an important prerequisite for prevention programmes to focus on increasing people's knowledge about sexual transmission, and, to overcome the misconceptions that act as a disincentive to behaviour change. Indicators on sexual behaviour and the promotion of safer sexual behaviour are at the core of AIDS programmes, particularly with young people who are not yet sexually active or are embarking on their sexual lives, and who are more amenable to behavioural change than adults. Finally, higher risk male-male sex reports on unprotected anal intercourse, the highest risk behaviour for HIV among men who have sex with men.

Knowledge of HIV prevention methods

Prevention indicator: Percentage of young people 15-24 who both correctly identify two ways of preventing the sexual transmission of HIV and who reject three misconceptions about HIV transmission.

Year	Male	Female

Comments:

Source:

Reported condom use at last higher risk sex (young people 15-24)

Prevention indicator: Proportion of young people reporting the use of a condom during sex with a non-regular partner.

Year	Male	Female

Comments: For this indicator only data will be shown if they were collected after 1998.

Source:

Age-mixing in sexual partnerships among young women

The proportion of young women who have had sex in the last 12 months with a partner who is 10 or more years older than themselves.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:

Source:

Reported non-regular sexual partnerships

Prevention indicator: Proportion of young people 15-24 having at least one sex partner other than a regular partner in the last 12 months.

Year	Male	Female

Comments:

Source:

Knowledge and behaviour (continued)

Ever used a condom

Percentage of people who ever used a condom.

Year	Area	Age group	Male	Female	All
------	------	-----------	------	--------	-----

Comments:
Source:

Adolescent pregnancy

Percentage of teenagers 15-19 who are mothers or pregnant with their first child.

Year	Percentage
------	------------

Comments:
Source:

Age at first sexual experience

Proportion of 15-19 year olds who have had sex before age 15.

Year	Male	Female

Comments:
Source:

Prevention indicators

Male and female condoms are the only technology available that can prevent sexual transmission of HIV and other STIs. Persons exposing themselves to the risk of sexual transmission of HIV should have consistent access to high quality condoms. AIDS Programs implement activities to increase both availability of and access to condoms. These activities should be monitored and have resources directed to problem areas. The indicator below highlights the availability of condoms. However, even if condoms are widely available, this does not mean that individuals can or do access them.

Condom availability nationwide

Total number of condoms available for distribution nationwide during the preceding 12 months, divided by the total population aged 15-49.

Year	N	Rate
------	---	------

Comments:
Source:

Prevention of mother-to-child transmission (MTCT) nationwide

Percentage of women who were counselled during antenatal care for their most recent pregnancy, accepted an offer of testing and received their test results, of all women who were pregnant at any time in the preceding two years.

Year	N	Rate
------	---	------

Comments:
Source:

Blood safety programs aim to ensure that the majority of blood units are screened for HIV and other infectious agents. This indicator gives an idea of the overall percentage of blood units that have been screened to high enough standards that they can confidently be declared free of HIV.

Screening of blood transfusions nationwide

Percentage of blood units transfused in the last 12 months that have been adequately screened for HIV according to national or WHO guidelines.

Year	N	Rate
------	---	------

Comments:
Source:

Sources

Data presented in this Epidemiological Fact Sheet come from several sources, including global, regional and country reports, published documents and articles, posters and presentations at international conferences, and estimates produced by UNAIDS, WHO and other United Nations agencies. This section contains a list of the more relevant sources used for the preparation of the Fact Sheet. Where available, it also lists selected national Web sites where additional information on HIV/AIDS and STI are presented and regularly updated. However, UNAIDS and WHO do not warrant that the information in these sites is complete and correct and shall not be liable whatsoever for any damages incurred as a result of their use.

Cheng, H., J. P. Zhang, S. D. Zhao, et al. 1994 Epidemiological Pattern of HIV Infection in Yunnan Province, China Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Session 042C.

Choi, K. H., C. Tian, X. Zheng, et al. 1994 Are STD Patients at Risk for HIV Infection in China? Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Poster P.C.0067.

Choi, K. H., X. Zheng, H. Zhou, et al. 1999 Treatment Delay and Reliance on Private Physicians among Patients with Sexually Transmitted Diseases in China International Journal of STD and AIDS, vol. 10, no. 5, pp. 309-315.

China Ministry of Health 1999 UNAIDS Country Profile Update: April 1999 Report

China National AIDS Program 1999 Update UNAIDS Epidemiological Fact Sheet UNAIDS.

Chen, J., S. Lai, W. Liu, et al. 2000 Epidemiological Characteristics of HIV Infection among IDUs in Guangxi, China 1996-1999 XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract MoPeC2400.

Chen, X. S., Y. P. Yin, G. J. Liang, et al. 2001 Prevalence and Risk Factors for Gonococcal and HIV Infections among Female Sex Workers in China Presented at International Congress of Sexually Transmitted Infections, ISSTD/ISTD, Berlin, Germany, 24-27 June 2001, Abstract in International Journal of STD & AIDS, vol. 12, suppl. 2, pp. 179-180.

Chen, Z. D., X. H. Liu, B. Yu, et al. 2002 HIV Infection and Condom Use among Sex Workers in Wuhan, China XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract MoPeC3490.

Hehe, C., Z. Jiapeng, P. Songfeng, et al. 2000 Analysis and Prediction on the Trends of HIV Infection Epidemic in Yunnan Province Journal of China AIDS/STD Prevention and Control, vol. 6, no. 5, pp. 257-260.

Huifen, Y., A. Xiaojing, J. Manhong, et al. 2001 Report on HIV/AIDS Surveillance in Yunnan Province in 1994 Journal of China AIDS/STD Prevention and Control, vol. 7, no. 2, pp. 74-76.

Liao, S. S. 1997 HIV/AIDS in China Presented at Monitoring the AIDS Pandemic (MAP) in Asia Symposium, Network Consultative Meeting, 10/21-23, Manila, Philippines.

Laeyendecker, O. 2002 Molecular Epidemiology of HIV-1 Circulating Recombinant Forms CRF01_AE and CRF08_BC in Guangxi Province, China 9th Conference on Retroviruses and Opportunistic Infections, Foundation for Retrovirology and Human Health, Seattle, WA., 2/24-28, online abstract, Poster Session 100, #761-W.

Li, L., Z. Guoying, M. Xiaoying, et al. 2000 Summary of HIV Sentinel Surveillance for 1995 to 1999 in Sichuan Province Journal of China AIDS/STD Prevention and Control, vol. 6, no. 5, pp. 280-281.

Mao, L., L. Peng, L. Hui, et al. 2001 Epidemiological Study on HIV/AIDS in Guangdong Province Journal of China AIDS/STD Prevention and Control, vol. 7, no. 1, pp. 11-13.

Qu, S., X. H. Sun, X. Zheng, et al. 1998 HIV Sentinel Surveillance in China in 1997 12th World AIDS Conference, Geneva, 6/28 - 7/3, Poster 13145.

Qu, S., X. H. Sun, X. Zheng, et al. 1996 Establishment of HIV Sentinel Surveillance System in China and Report of Surveillance in 1995 Chinese Journal for STD and AIDS Prevention and Control, vol. 2, no. 5, pp. 193-197.

Qu, S., X. H. Sun, X. Zheng, et al. 1997 The Report of HIV Sentinel Surveillance in China: 1996 Chinese Journal for STD and AIDS Prevention and Control, vol. 3, no. 5, pp. 193-197.

Qu, S., X. H. Sun, X. Zheng, et al. 1998 HIV Sentinel Surveillance in China in 1997 Chinese Journal for STD and AIDS Prevention and Control, vol. 4, no. 4, pp. 178-185.

Qu, S., X. Sun, X. Zheng 2000 National Sentinel Surveillance of HIV Infection in China from 1995 to 1998 XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster MoPeC2379.

Qu, S., et al. 2001 National HIV/AIDS Sentinel Surveillance Report in China, 2000 Department of Diseases Control, Ministry of Health, National Center for AIDS Prevention and Control, National Coordination Group for HIV Sentinel Surveillance, unpublished report.

Wei, L., J. Chen. R. Li, et al. 2000 Peer Education for HIV/STD Prevention among Community-Based Drug Users in Guangxi Province, Southern China XIII International AIDS Conference, Durban, South Africa, 7/9-14, Abstract WeOrC500.

Xinhua, S., N. Junhua, G. Qili 1994 AIDS and HIV Infection in China AIDS, vol. 8, suppl. 2, pp. S55-S59.

Zhang, L., G. M. Qing, G. Liu, et al. 1993 HIV/AIDS Trends in Sichuan Province IX International Conference on AIDS, Berlin, 6/6-11, Abstract PO-C08-2781.

Zheng, X. W., J. P. Zhang, C. Q. Tian, et al. 1993 Cohort Study of HIV Infection among Drug Users in Ruili, Longchuan and Luxi of Yunnan Province, China Biomedical and Environmental Sciences, vol. 6, pp. 348-351.

Zhang, J. P., H. H. Cheng, S. D. Zhao, et al. 1994 Research on the Epidemic of HIV Infection in Yunnan Province China Tenth International Conference on AIDS, Yokohama, Japan, 8/7-12, Abstract P.C.0066.

Zheng, X., J. Zhang, X. Qu., et al. 1996 The Sero-Prevalence and Incidence Rates of HIV Infection among IDU by Cohort Study from 1992 to 1995 in Ruili and Other ... XI International Conference on AIDS, Vancouver, 7/7-14, Abstract Tu.C.2515.

Zhang, J., B. Xiuqiong, C. Hehe, et al. 1998 HIV Prevalence Trends among Intravenous Drug Users in Yunnan Province between 1992-1997 12th World AIDS Conference, Geneva, 6/28 - 7/3, Poster 43459.

Zheng, X., Z. Jiapeng, Q. Shuquan, et al. 1998 A Cohort of HIV Infection among IDU's and Natural History of HIV Infection in Ruili, China, 1992-1997 12th World AIDS Conference, Geneva, 6/28 - 7/3, Poster 23244.

Zheng, X. 1997 HIV/AIDS in China Workshop on HIV, AIDS, and STD Epidemiology in Western Pacific Region, Manila, Philippines, 9/8-10, document no. 1.

Zhang, Y., F. Sun, M. Zhou, et al. 2000 HIV Testing among Pregnant Women in Xinjiang, China XIII International AIDS Conference, Durban, South Africa, 7/9-14, Poster MoPeC2424.

Zhu, O. Y., W. Liu, J. Chen, et al. 2002 Very Rapid Spread of HIV Infection among Persons at Risk in Guangxi, China, 1996-2000 XIV International AIDS Conference, Barcelona, Spain, 7/7-12, Abstract TuPpC2055.

Websites: **Joint United Nations Programme on HIV/AIDS:** <http://www.unchina.org/un aids/>

Annex: HIV surveillance by site

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
Pregnant women	Major urban areas	Guangdong Province											0	0				
		Guangxi Province (1), Guangxi Provi									0	0	0	0	0			
		Guangxi Province (2), Guangxi Provi													0			
	Outside major urban areas	Baoshan											0	0.54				
		Dali											0	1.63				
		Kaiyuan											0.47	0.14				
		Kunming											0.11	0.42				
		Longchuan											0.39	0.14				
		Quijing											0	0.12				
		Sichuan Province													0			
		Ximao											0.22	0				
		Xinjiang Province														0.50		
		Yingjiang											0.12	0.12				
		Yining										0	0.40	0.88				
		Yining Women's & Children's Hospita												0.62				
		Yunnan Province				0.20	0	0.04						0.22				
Sex workers	Major urban areas	Anhui Province								0	0.19	0.22	0.50	0	0.19			
		Beijing								0		0.15	0	0	0.20			
		Fujian Province								0	0	0	0	0	0			
		Fushun													0			
		Guangdong Province									0	0	0.14	0.32				
		Guangxi Province (1), Guangxi Provi									0	0	2.55	6.19	10.29			
		Guangzhou Province								0	0	0	0.30	0.80	2.20			
		Guizhou Province								0	0	0	0	0.50	0.19			
		Henan Province								0	0	0.62	0	0	0			
		Hubei Province								0	0	0.19	0	0	0			
		Hunan Province								0	0	0	0	0.16	0.25			
		Jiangsu Province													0			
		Kunming									0.71			2.24			10.30	
		Liaoning Province								0	0	0	0	0.49				
		Masanjia													0			
		Mianyang											0	0	0			
		Nanning								0		0.70	2.25					
		Shaanxi Province								0	0	0	0	0	0			
		Shandong Province								0	0	0	0	0	0			
		Shanghai Province								0	0	0						
	Outside major urban areas	Shanxi Province								0			0	0	0			
		Shenzhen										0	0	0	0			
		Tianjin Province								0	0	0	0	0.20	0			
		Zigong											0	0				
		Chengdu								0.10	0	0	0.20	0.40	0			
		Hainan Province								0.17	0	0	0	0	0			
		Wuhan City										0.18		0	0.24		0.44	
		Xinjiang Province											23.08	0.75				
		Yunnan Province			0	0	0.38	0	0	0	0.50	1.50	2.67	0.90	2.70	2.90		
Injecting drug users	Major urban areas	Beijing													0			
		Changsha													0.59			
		ChongQing Province											0	0.44	0.82			
		Fujian Province												0	0			
		Gansu Province				0				0	0	0	0	1.88				
		Guangdong Province								0	0.22	1.05	3.85	21.85	20.46			
		Guangxi Province (1), Guangxi Provi								0	0	2.26	10.96	12.94	16.45			
		Guiyang											1.56	2.94	0.66			
		Guizhou Province								0	0.17	0.15						
		Hubei Province										0.18	0	0	0			
		Hunan Province											0.33	0.80				
		Inner Mongolia Province				0	0			0	0	0.18						
		Jiangsu Province													0.29			

Group	Area	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
Injecting drug users	Major urban areas	Jiangxi Province											0	0.15	8.95			
		Kunming					0	0	0	0.68	9.29	27.52	23.40	24.40				
		Liaoning Province													0			
		Liuzhou									0	2.00	7.00					
		Nanning									0		6.30					
		Pan'shi											7.06					
		Panzhihua												1.11	4.29			
		Shaanxi Province								0	0	0			0			
		Shanxi Province														0		
		Sichuan Province								0	0	4.15						
		Urumqi											40.45					
		Zhejiang Province													0			
	Outside major urban areas	Baoshan					0	0	0.63	0.45	2.36	9.64	5.90	6.80				
		Binchuan									51.79							
		Binyang													18.84			
		Binyang county									0	0		17.00				
		Cangling									23.14		13.80	17.60				
		Changlu															0.42	
		Dali					0	0	0	4.66	47.50		57.40	48.90				
		Dehong											63.40	69.80				
		Dujun											1.69	1.04	0			
		Geju									8.62							
		Gengma									49.64							
		Jinghong											1.71	0.89				
		Kaiyuan					0	0	0	0.38	25.67	70.27	70.60	58.20				
		Lancang								2.32	27.87							
		Lijiang													2.60			
		Lincang county							4.86	50.00	75.00	66.67	52.90	57.10				
		Longchuan					42.96	40.00	39.66	43.20	45.45							
		Longchuan county						46.03	47.13	60.00		56.67						
		Luxi					3.73				15.25							
		Luxi county						0	1.54	8.27			20.90	47.10				
		Ningxia Province											2.15	0.03	0.70			
		Pingxiang									16.00	22.00	19.00		15.38			
		Qinghai Province										0	0.55	0.29	0.53			
		Ruli		45.14	67.98		81.54	36.54		50.00	76.79							
		Ruli county						56.25		73.21								
		Simao											7.90	13.20				
		Tiandong county									0	0						
		Tianyang county									0	0	23.00					
		Weishan									12.24							
		Weishan county							0			59.52						
		Wenshanpanlong											65.00	75.00				
		Wulumuqi													37.92			
		Wushi															38.68	
		Xichang											6.67	0	3.05			
		Xinjiang Province								0.15	12.62	24.12						
		Yingjiang county					17.44		35.29									
		Yining											85.12	81.77	82.49			
		Yunnan Province		38.81	26.64	3.73	1.67	2.67					24.40	27.82				
		Yuxi									16.67		13.70	12.90				
		Zhaotong city											5.70	3.50				
STI patients	Major urban areas	Anhui Province								0	0	0	0	0.10	0			
		Beijing						0					0					
		Changzhou											0	0.13	0			
		Dalian								0	0	0	0	0	0			
		Dandong											0	0	0			
		Fujian Province								0.10	0	0	0.40	0	0			
		Guangdong Province								0	0	0.03	0.03	0.04				
		Guangxi Province (1), Guangxi Provi									0	0	0.27	0.30	0.36			
		Guangzhou Province								0	0	0.12	0.20	0	0.38			
		Guizhou Province												0	0			
		Hangzhou								0	0	0	0					
		Hebei Province										0	0					
		Heilongjiang Province											0	0	0			
		Henan Province								0	0	0.19	0	0.77	0.76			
		Hubei Province								0	0	0	0	0	0			

Group	Area		1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
STI patients	Major urban areas	Hunan Province									0	0	0	0	0	0				
		Inner Mongolia Province													0	0				
		Jiangsu Province										0	0.12	0.13		0				
		Jiangxi Province										0	0	0	0	0	0			
		Jilin Province												0	0	0	0			
		Jinan										0	0	0	0					
		Kunming											0.45		0.26	0.45				
		Liaoning Province										0	0	0						
		Liuzhou													0	0.47	1.33			
		Maoming														0.16	1.01			
		Nanjing																0		
		Nanning										0	0	0	0.40	0.16	0			
		Ningbo													0	0	0			
		Qingdao																0		
		Qinghai Province																0		
		Shaanxi Province																0		
		Shandong Province														0				
		Shanghai Province											0	0	0	0	0	0		
		Shanxi Province													0	0	0	0		
		Shenzhen										0			0	0	0	0		
		Shenzhen Province													0		0	0		
		Shijazhuang														0	0			
		Sichuan Province								0			0	0	0	0	0	0		
		Suzhou											0	0.13	0.25	0.15	0.14	0.09		
		Tianjin Province											0	0	0	0	0	0		
		Zhejiang Province											0	0						
		Zhongshan															0.39	0		
	Outside major urban areas	Baise													0.30	0.24	0			
		Baoshan										6.70	1.03		0.96	4.06				
		Binchuan											15.26		1.40	0.55				
		Chengdu										0.03				0.08				
		Chuxiong											0.83		0	0.36				
		Dandong																0		
		Dehong													2.10	0.95				
		Gejiu											0.23		2.38	6.29				
		Gengma														0				
		Hainan Province										0	0	0	0	0	0			
		Jinghong													0	0				
		Lancang											0.67		5.08	11.32				
		Lijiang													11.27	6.25				
		Luxi									1.40	0	1.44							
		Pingxian Province												0						
		Pingxiang													0	0.30	0.46	0.15		
		Quijing														0.38	0			
		Ruili										1.90	2.25		6.47	5.94				
		Wenshan													0.43	2.27				
		Yunnan Province					0	0	0					0	0.40	0.20	0.30			
		Zhaotong														0.44	0.53			
Men having sex with men																				
Tuberculosis patients																				